

ERRATA

One-Loop Threshold Effects in String Unification[★]Vadim S. Kaplunovsky[†]*Department of Physics, University of Texas, Austin, TX 78712, USA[‡]*

As originally published, the article contained a few minor errors. Unfortunately, these errors have lately caused a not-so-minor confusion among several physicists working on the phenomenology of string unification. To avoid further confusion, I would like to make the following corrections:

(1) The renormalization scheme referred throughout the article as $\overline{\text{MS}}$ is in fact the modified minimal subtraction scheme for the *dimensional reduction* and not the standard dimensional regularization. The proper name for the scheme I used is $\overline{\text{DR}}$.

(2) In the second paragraph on page 154 the formula for ξ'' should be $\xi'' = 1 + \log(2/\sqrt{27}\pi) - \gamma \approx -1.6767$ instead of $\xi'' = 1 - \log(2/\sqrt{27}\pi) - \gamma \approx -0.532$.

The rest of the mistakes involve misplaced factors of 2:

(3) Formula (26) for the effective scale of string unification should read

$$M_{\text{GUT}} \stackrel{\text{def}}{=} \frac{2e^{(1-\gamma)/2} 3^{-3/4}}{\sqrt{2\pi\alpha'}} \approx \frac{e^{(1-\gamma)/2} 3^{-3/4}}{4\pi} g_{\text{string}} M_{\text{Planck}} \approx g_{\text{GUT}} \times 5.27 \cdot 10^{17} \text{ GeV}. \quad (26)$$

(In the original publication, the factor **2** shown here in the boldface was missing.) Note that the correct formula here uses the tree-level relation $k_a g_a^2 = g_{\text{GUT}}^2 = 32\pi/\alpha' M_{\text{Planck}}^2$ which differs by a factor of 2 from a similar formula given in ref. [6]. This difference is due to different normalization conventions for the gauge generators Q_a and gauge couplings

★ Originally published in *Nucl. Phys.* **B307** (1988), pp. 145–156

† Current research supported in part by the NSF under grant #PHY-90-09850 and by the Robert A. Welch Foundation. Research leading to the original publication was supported in part by the NSF under grant #PHY-86-12280

‡ Current address.

g_a — unlike Ginsparg, I used the phenomenological convention[§] throughout the paper and adjusted the string-theoretical formulæ to work with this convention.

(4) In formulæ (1), (2), (7) and (24) coefficients printed as $4\pi^2$ should be $16\pi^2$. Similarly, in formulæ (5) and (21) coefficients in front of the respective integrals should be $1/16\pi^2$ instead of $1/4\pi^2$ and in formula (16) the coefficient printed as $2\alpha'^2$ should be $\alpha'^2/2$.

(5) In formulæ (6) and (23), the expressions for the coefficients b_a of the low-energy β -functions were too small by a factor of 2. Correspondingly, the field-theoretical functions $\mathbf{B}_a(t)$ and the string-theoretical functions $\mathcal{B}_a(\tau, \bar{\tau})$ also missed that factor. The correct form of the field-theoretical eq. (5) is

$$\mathcal{W}_a^{\text{field}} = \frac{1}{16\pi^2} \int_0^\infty \frac{dt}{t} C_\Lambda(t) \cdot \left[\mathbf{B}_a(t) \stackrel{\text{def}}{=} \mathbf{2} \text{str} \left(Q_a^2 \left(\frac{1}{12} - \chi^2 \right) e^{-tM^2} \right) \right]. \quad (5)$$

Consequently, the super-traces in eqs. (7) and (8) should also be multiplied by two; in particular, the correct formula for the one-loop threshold corrections in GUTs is

$$\Delta_a = \mathbf{2} \text{str}_{M \sim M_{\text{GUT}}} \left(Q_a^2 \left(\frac{1}{12} - \chi^2 \right) \log \frac{M_{\text{GUT}}^2}{M^2} \right). \quad (8)$$

Similarly, the correct form of the string-theoretical formula (22) is

$$\mathcal{B}_a(\tau, \bar{\tau}) = \frac{\mathbf{2}}{|\eta(\tau)|^4} \sum_{\text{even } \mathbf{s}} (-)^{s_1+s_2} \frac{dZ_\Psi(\bar{\tau}, \mathbf{s})}{2\pi i d\bar{\tau}} \cdot \text{Tr}_{s_1} \left(Q_a^2 \cdot (-)^{s_2 F} q^H \bar{q}^{\bar{H}} \right)_{\text{int}}. \quad (22)$$

The relation (25) between these $\mathcal{B}_a$ functions and the

§ Most phenomenologists normalize the non-abelian gauge generators to $\text{tr}(Q_a^2) = \frac{1}{2}$ where the trace is taken over the fundamental representation of an $SU(N)$ group such as $SU(3)_{\text{color}}$ or $SU(2)_{\text{weak}}$; correspondingly, the generators of a GUT group are normalized to $\text{tr}_{\mathbf{5}}(Q_a^2) = \frac{1}{2}$ for the $SU(5)$ or $\text{tr}_{\mathbf{10}}(Q_a^2) = 1$ for the $SO(10)$. On the other hand, many string theorists normalize the generators to $\text{tr}(Q_a^2) = 2$ where the trace is taken over a vector representation of an $SO(2N)$ group such as $SO(32)$ or $SO(16) \subset E_8$; according to this convention, phenomenologically-normalized Q_a should be multiplied by $\sqrt{2}$. In both conventions, the gauge connection A_μ is $igQ_a A_\mu^a$, A_μ^a being canonically-normalized vector fields; therefore, the string-theoretical convention has to compensate for the Q_a being bigger by a factor $\sqrt{2}$ by having the gauge couplings g being smaller by the same factor.

string threshold corrections Δ_a remains unchanged, which means that the actual values of Δ_a should be doubled.

As to the formulæ (6) and (23) for b_a and the infra-red limits of $\mathbf{B}_a(t)$ and $\mathcal{B}_a(\tau, \bar{\tau})$, in addition to missing an overall factor of 2, they also used inconsistent conventions for the traces. The correct formulæ should read

$$\mathbf{B}_a(t) \xrightarrow{t \rightarrow \infty} -\frac{11}{3} \text{tr}_{V,M=0}(Q_a^2) + \frac{2}{3} \text{tr}_{F,M=0}(Q_a^2) + \frac{1}{3} \text{tr}_{S,M=0}(Q_a^2) \equiv b_a, \quad (6)$$

$$\mathcal{B}_a \xrightarrow{\tau_2 \rightarrow \infty} -\frac{11}{3} \text{tr}_{V,M=0}(Q_a^2) + \frac{2}{3} \text{tr}_{F,M=0}(Q_a^2) + \frac{1}{3} \text{tr}_{S,M=0}(Q_a^2) = b_a \equiv \lim_{t \rightarrow \infty} \mathbf{B}_a(t), \quad (23)$$

where the traces are taken over the massless charged particles *and count each CPT-conjugate particle-antiparticle pair only once*.

Most of the above errors were either obvious or irrelevant to most of the readers of the paper. Unfortunately, the missing factor of 2 in eqs. (22) and (23) was neither, which lead to its propagation via papers concerned with the threshold corrections in specific string models. I would like to use this opportunity to apologize to the authors of those papers for leading them into the error. I would also like to apologize for propagating this error myself, in the article “*Moduli-Dependence of String Loop Corrections to Gauge Coupling Constants*” I co-authored with L. Dixon and J. Louis (*Nuclear Physics* **B355** (1991), p. 649). Fortunately, only some of the intermediate results of that article are affected by this missing factor of 2; the final results — the relations between Δ_a of an orbifold and the b'_a (b_a coefficients of the $N = 2$ partial orbifold) are correct, provided one uses the correct definitions of the b_a and b'_a coefficients.

The author thanks Ignatios Antoniadis, for pointing out the above errors (even though his attempts to correct them were also erroneous), and Mirjam Cvetič, for convincing me that these errors must be corrected and for helping me to correct some of them.